



Science Mission Directorate
National Aeronautics and Space Administration

UAH

An Update on Regional Profile Assimilation and Near-Real-Time Modeling Plans

Brad Zavodsky

University of Alabama in Huntsville
Huntsville, Alabama

Shih-hung Chou, Gary Jedlovec, & Bill Lapenta

NASA / Marshall Space Flight Center
Huntsville, Alabama

AIRS Science Team Meeting – September 2006



transitioning unique NASA data and research technologies to the NWS



Outline

- Motivation
- Overview of Previous AIRS Science Team Meeting Presentations
- New Results from Assimilation/Forecasting Work
- Near-Real-Time (NRT) AIRS Assimilation
- Brief Discussion






Motivation

- AIRS data complements traditional upper-air observations in data sparse regions (e.g. ocean).
- Hyperspectral nature of AIRS sounder allows for highest vertical resolution of any current remote sensing system
- In-house computing resources at SPoRT are currently not able to handle radiance assimilation
- Level-II profiles provide a straightforward method for obtaining information in data-void regions without running complex RTA
- Profiles are easier to handle than radiances in a NRT environment

Sample swath of AIRS data with prototype V5.0 quality indicators





Summary of Recent Results


- ADAS configured to optimally assimilate AIRS thermodynamic profiles intelligently using quality indicators (QIs) to determine largest volume of highest quality data
- Short WRF forecast used as background for ADAS analysis (previously initialized with GFS, but discussing moving to the NAM), WRF produced 65-h forecasts
- Two case studies: west coast (V4.0, Jan. 2004) and east coast (prototype V5.0, Nov. 2005)
- Impact in temperature and moisture analyses in region of AIRS data
- Overall positive impact on temperature and moisture forecasts with the inclusion of AIRS profiles
- Prototype V5.0 data reduced bias and RMSE of temperature and moisture in forecasts when compared to RAOBs—improvements in RTA and QC






Case Study: November 20-22, 2005


WRF Domain for November 2005 Case Study




- WRF domain covers most of CONUS—similar domain to be used for testing NRT
- Rapidly intensifying storm in region relevant to SPoRT interests
- Under forecasted by operational models at storm's strongest point—potential impact
- Ample verification data available over the Eastern US (minimal terrain issues)



Surface analysis 11/20/05 12 UTC



Surface analysis 11/21/05 12 UTC



Surface analysis 11/22/05 12 UTC




transitioning unique NASA data and research technologies to the NWS



53h Forecast Impact—Mean Sea Level Pressure

Valid at 1200 UTC 22 November 2005



- Low pressure area deepens by more than 4 hPa with inclusion of AIRS data
- Storm center shifts slightly to the south and west
- Both alterations are improvements over the control when compared to surface analysis




transitioning unique NASA data and research technologies to the NWS



41h Forecast Impact—6h Cumulative Precipitation


Valid at 0000 UTC 22 November 2005





NRT Assimilation


- Single case studies are not necessarily representative (statistically significant) of overall model performance
- Twice daily model runs (AM and PM) for a contiguous period
- Initial sensitivity study using *over water* soundings (latest prototype V5.0) will help determine optimal data set for later NRT assimilation:
 - **CNTL:** control; use no AIRS data
 - **NOQC:** use all AIRS data without regard for QIs
 - **BEST:** use only full soundings
 - **ALLW:** use QIs to select the highest quality data
- Troubleshoot process, determine case studies, and provide results for a journal article
- Date, time, and location of AIRS data will control which data are used and will run system from pre- through postprocessing





NRT Assimilation

- NAM is run 4 times daily with a lag time of ≈ 2 hours; ≈ 3 hours for 0-48 hour forecasts
- 1-hr WRF forecasts can be completed in 30 seconds; 48-hr forecast can be completed in under 9 minutes using full capabilities of our cluster
- Preprocessing of AIRS profiles into ADAS format takes ≈ 5 minutes
- Assimilation of AIRS profiles in current configuration takes ≈ 10 minutes
- Time lag of AIRS data processing—how long after observation time until data are available? Ideally, we would like AIRS data to be available ≤ 2 hours after valid time*





Verification of Forecast Impact

- Verification at 6-hour intervals at each grid point using analyses (F00) collocated in time to calculate bias and RMSE for:
 - **temperature, specific humidity, height, MSLP, winds**
- Verification at 3-hour intervals in each grid box using extrapolated NCEP Stage IV precipitation data to calculate equitable threat scores and bias scores for:
 - **6h and 24h cumulative precipitation totals**
- Time series of MSLP for the forecast period for each case against METAR data along the east coast
- If month-long set of cases yields extra-tropical cyclone events, storm tracks between the various model runs and observations can be compared
- Results will be posted daily on the SPoRT website after forecast run to track the statistics





Conclusions/Wrap-Up

- Improvements in temperature, moisture, MSLP, and precipitation fields at various forecast times with addition of AIRS profiles for one case study
- More thorough assessment through extended NRT study
- AIRS Level-II thermodynamic profiles are desired in NRT with less than 2 hours lag for continued research
- We would like to test our system with at least a month of NRT prototype V5 data prior to the official V5 release





Science Mission Directorate
National Aeronautics and Space Administration

UAH

Supplemental Slides



transitioning unique NASA data and research technologies to the NWS

